

b.) Remarks

Claim 51 is objected to as containing a typographical error. Such formal matter has been attended to above, in conformity with the Examiner's kind suggestion.

Claims 3-4 and 50 are rejected under 35 U.S.C. 112, first paragraph, because the Examiner contends the specification does not reasonably enable one of ordinary skill to practice the scope of the invention. The Examiner states (page 7, line 18 to page 8, line 1) that without guidance, random screening is excessive.

At the outset, Applicants respectfully wish to point out this is not what *Ex parte Forman* says; rather the specification is enabling if it presents a reasonable amount of guidance or if random experimentation is routine in the art (230 USPQ at 547). In this regard, the Examiner has made no *prima facie* showing that the required level of experimentation, even if random, is not routine. Indeed, it plainly is where, as here, such can simply be automated. Nonetheless, solely in order to reduce the issues and expedite prosecution, Applicants have above cancelled these claims without prejudice or disclaimer. Accordingly, the rejection is mooted.

Claims 2-9, 11-13 and 48-52 are rejected under 35 U.S.C. 102(a) as being anticipated by Isshiki et al. (J. Biol. Sci. Vol, 274(18):12499-12507, Apr. 1999) or Zhou D. et al. (Eur. J. Biochem. Vol. 263(2):571-576). Claim 10 is rejected under 35 U.S.C. 103(a) as being unpatentable over Isshiki et al. or Zhou D. et al.

Initially, it is noted that Isshiki is dated "April 1999", which date follows the February 25, 1999 filing date of Applicants' Japanese priority application JP 99-47571. Additionally, Applicants' investigation has determined that Zhou was not published, at the

earliest, until July 11, 1999.<sup>1/</sup>

The rejections over both Isshiki and Zhou can be overcome by filing a sworn translation of Applicants' Japanese priority application. Accordingly, such translation is enclosed.

In view of the above amendments and remarks, and the accompanying sworn translation, Applicants submit that all of the Examiner's concerns are now overcome and the claims are now in allowable condition. Accordingly, reconsideration and allowance of this application is earnestly solicited.

Claims 2, 5-25, 29-31, 33, 38, 48, 49 and 51-54 remain presented for continued prosecution. Rejoinder of claims 14-25, 29-31, 33, 38, 53 and 54, each of which relate to a process of using (claims 18-25, 53 and 54), producing (claims 14-17) or determining (claims 29-31, 33 and 38) the elected subject matter, is respectfully requested

---

<sup>1/</sup> Applicants have learned Zhou was publicly available, according to the publisher, on July 12, 1999 (Tab A). It was first received in the Senckenberg library on July 26, 1999, and was not published online until December 25, 2001 (Tab B). The reference itself is dated July 11, 1999 (Tab C).

Applicants' undersigned attorney may be reached in our New York office by telephone at (212) 218-2100. All correspondence should continue to be directed to our below listed address.

Respectfully submitted,

A handwritten signature in black ink, appearing to read "Lawrence S. Perry", written over a horizontal line.

Attorney for Applicants  
Lawrence S. Perry  
Registration No. 31,865

FITZPATRICK, CELLA, HARPER & SCINTO  
30 Rockefeller Plaza  
New York, New York 10112-3801  
Facsimile: (212) 218-2200

LSP\ac

NY\_MAIN 424438v1

**Wolfgang Völger**

---

Von: Dobson Katherine [Katherine.Dobson@oxon.blackwellpublishing.com]  
Gesendet: Dienstag, 30. März 2004 10:39  
An: Dr. Wolfgang Völger  
Betreff: RE: publishing date (EJBIOCHEM Feedback Form)

Dear Dr Völger

The official publication date of this issue is 12 July 1999.

Best wishes

Katherine Dobson

-----Original Message-----

From: Dr. Wolfgang Völger [mailto:voelgerw@tbk-patent.de]  
Sent: 10 March 2004 09:56  
To: ejbiochem-feedback@highwire.stanford.edu  
Cc: voelgerw@tbk-patent.de  
Subject: publishing date (EJBIOCHEM Feedback Form)

---

Comments sent via EJB Feedback Page

---

NAME: Dr. Wolfgang Völger  
EMAIL: voelgerw@tbk-patent.de  
IP ADDRESS: 212.202.136.114  
HOSTNAME: port-212-202-136-114.reverse.qsc.de  
PREVIOUS PAGE: <http://www.ejbiochem.org/content/vol263/issue2/index.shtml>  
BROWSER: Mozilla/4.0 (compatible; MSIE 6.0; Windows NT 5.0)  
PROMOTIONAL USE: Not granted

---

COMMENTS:

Dear Madame and Sir,

I am looking for the exact publishing date of

European Journal of Biochemistry, Volume 263, Issue 2,  
July 1999

e.g. when it was actually delivered to the public in  
US and Europe. Could you please give me this  
information including a short confirmation for our  
clients...

Thank you very much in advance.

Sincerely yours,

Dr. Wolfgang Völger  
TBK-Patent  
Bavariaring 4-6  
80336 München - Germany

**Blackwell Synergy**

Home Browse Search MySynergy Register Help

Username:  [Zitrona Login](#)

Password:  [Login](#)


[Forgotten Password?](#) [Logout](#)

You are at: [Home](#) > [List of Issues](#)

## List of Issues

### European Journal of Biochemistry

Published on behalf of the Federation of European Biochemical Societies



Go to:  [Go](#)

**\*\*Sign-up for FREE email table of contents alerts\*\***  
**\*\*Add this journal to your favorites\*\***

Submit your manuscript to The FEBS Journal online at [Manuscript Central](#). Click here for full [Author Guidelines](#).

**\*Free - all review articles published to date in EJB\***

OnlineEarly (Peer-reviewed full-text articles for future issues)

Review articles

☐ 2004 271(1) - 271(6)

- March 2004 (Vol. 271 Issue 6 Page 1065-1227) **Latest Issue**
- March 2004 (Vol. 271 Issue 5 Page 877-1064)
- February 2004 (Vol. 271 Issue 4 Page 663-875)
- February 2004 (Vol. 271 Issue 3 Page 459-661)
- January 2004 (Vol. 271 Issue 2 Page 237-458)
- January 2004 (Vol. 271 Issue 1 Page 1-235)

☐ 2003 270(1) - 270(24)

☐ 2002 269(1) - 269(24)

☐ 2001 268(1) - 268(24)

☐ 2000 267(1) - 267(24)

☐ 1999 259(1-2) - 266(3)

- August 1999 (Vol. 263 Issue 3 Page 605-927)
- July 1999 (Vol. 263 Issue 2 Page 295-603)
- July 1999 (Vol. 263 Issue 1 Page 1-291)
- June 1999 (Vol. 262 Issue 3 Page 627-947)
- June 1999 (Vol. 262 Issue 2 Page 247-624)
- May 1999 (Vol. 262 Issue 1 Page 1-245)
- May 1999 (Vol. 261 Issue 3 Page 593-821)
- April 1999 (Vol. 261 Issue 2 Page 347-590)
- March 1999 (Vol. 261 Issue 1 Page 1-344)
- March 1999 (Vol. 260 Issue 3 Page 571-927)
- March 1999 (Vol. 260 Issue 2 Page 293-S1)
- February 1999 (Vol. 260 Issue 1 Page 1-290)
- February 1999 (Vol. 259 Issue 3 Page 555-961)
- January 1999 (Vol. 259 Issue 1-2 Page 0-552)

☐ 1998 251(1-2) - 258(3)

QuickSearch :

☒ This Journal

☐ All Journals

[Search](#)

Further Search Options:

[Simple](#) [Advanced](#) [Saved](#)

Quicklink:

Vol:  Issue:  Page:

[Go](#)



Visit the [Journal Homepage](#) for more information on:

- How to subscribe
- Submitting a paper
- The journal's aims and scope
- The Editorial Board



- 552 **Post-translational modifications of the insect sulfakinins. Sulfation, pyroglutamate-formation and O-methylation of glutamic acid**

Reinhard Predel, Wolf Brandt, Roland Kellner, Jürgen Rapus, Ronald J. Nachman, Gerd Gäde

Online publication date: 25-Dec-2001

[Abstract](#)[References](#)[Full Text Article](#)[PDF \[845KB\]](#)

Highlight: ☐

- 561 **Expression of a higher plant light-harvesting chlorophyll *a/b*-binding protein in *Synechocystis* sp. PCC 6803**

Qingfang He, Thomas Schlich, Harald Paulsen, Wim Vermaas

Online publication date: 25-Dec-2001

[Abstract](#)[References](#)[Full Text Article](#)[PDF \[661KB\]](#)

Highlight: ☐

- 571 **Molecular cloning of a human UDP-galactose:GlcNAc $\beta$ 1,3GalNAc  $\beta$ 1,3 galactosyltransferase gene encoding an O-linked core3-elongation enzyme**

Dapeng Zhou, Eric G. Berger, Thierry Hennet

Online publication date: 25-Dec-2001

[Abstract](#)[References](#)[Full Text Article](#)[PDF \[673KB\]](#)

Highlight: ☐

- 577 **Engineering of conformations of plasminogen activator inhibitor-1. A crucial role of  $\beta$ -strand 5A residues in the transition of active form to latent and substrate forms**

Tove Kirkegaard, Signe Jensen, Susanne L. Schousboe, Helle H. Petersen, Rikke Egelund, Peter A. Andreasen, Kees W. Rodenburg

Online publication date: 25-Dec-2001

[Abstract](#)[References](#)[Full Text Article](#)[PDF \[686KB\]](#)

Highlight: ☐

- 587 **Chemical and antigenic structure of the O-polysaccharide of the lipopolysaccharides from two *Acinetobacter haemolyticus* strains differing only in the anomeric configuration of one glycosyl residue in their O-antigens**

Ralph Pantophlet, Simon R. Haseley, Evgeny V. Vinogradov, Lore Brade, Otto Holst, Helmut Brade


Online publication date: 25-Dec-2001

[Abstract](#)[References](#)[Full Text Article](#)[PDF \[1034KB\]](#)

Highlight: ☐

- 596 **Dynamin and Rab5 regulate GRK2-dependent internalization of dopamine D2 receptors**

[Abstract](#)[References](#)



[Home](#)
[Browse](#)
[Search](#)
[My Synergy](#)
[Register](#)
[Help](#)

Username:   
Password:   
[Forgotten Password?](#)

[Ziterra Login](#)  
[Login](#)  
[Logout](#)

You are at: [Home](#) > [List of Issues](#) > [Table of Contents](#) > [Full Text](#)

## Full Article

 [View/Print PDF article \(673K\)](#)

[Download to reference manager](#)

European Journal of Biochemistry

Volume 263 Issue 2 Page 571 - July (II) 1999

doi:10.1046/j.1432-1327.1999.00541.x

# Molecular cloning of a human UDP-galactose:GlcNAc $\beta$ 1,3GalNAc $\beta$ 1,3 galactosyltransferase gene encoding an O-linked core3-elongation enzyme

Dapeng Zhou, Eric G. Berger and Thierry Hennet

Using the full-length amino-acid sequences of the human  $\beta$ 1,3 galactosyltransferase ( $\beta$ 3GalT)-I, -II and -III enzymes as query, we have identified an additional member of the  $\beta$ 3GalT gene family within a sequenced region of the human chromosome 21 as found in GenBank. The novel human  $\beta$ 3GalT-V gene included an open reading frame of 933 bp encoding a protein of 310 amino acids with a short N-terminal cytoplasmic tail, a single predicted transmembrane domain and a large luminal catalytic domain. The human  $\beta$ 3GalT-V protein showed 34%, 27%, 31% and 23% sequence identity with the human  $\beta$ 3GalT-I, -II, -III and -IV enzymes, respectively. The expression of  $\beta$ 3GalT-V as a recombinant protein in Sf9 insect cells confirmed the galactosyltransferase activity catalyzed by this enzyme. Similarly to  $\beta$ 3GalT-I, -II and -III, the  $\beta$ 3GalT-V enzyme used  $\beta$ -linked GlcNAc as an acceptor, but unlike the former enzymes  $\beta$ 3GalT-V exhibited a marked preference for the O-linked core3 GlcNAc $\beta$ 1,3GalNAc substrate. The  $\beta$ 3GalT-V gene was mainly expressed in human small intestine and to a lesser extent in pancreas and testis. Although  $\beta$ 3GalT-V transcripts were not detected in normal colon tissue, based on Northern analysis,  $\beta$ 3GalT-V mRNA was found in the adenocarcinoma cell line Colo 205.

The availability of large DNA sequence databases has enabled the retrieval of structurally related genes based on similarity searches. This approach has been fruitful in glycobiology, as many glycosyltransferase activities previously believed to be encoded by single genes have turned out to represent families including several enzymes. Characterization of these families has revealed the complex pattern of tissue expression of the corresponding genes and the specificity of each protein towards distinct acceptor substrates, as exemplified by fucosyltransferase [ 1] and polypeptide N-acetylgalactosaminyltransferase enzymes [ 2].

In vertebrates, galactose is transferred either as  $\alpha$ - or  $\beta$ -anomer through 1,3 or 1,4 linkage (reviewed in [ 3]). Multiple  $\beta$ 1,4 galactosyltransferase ( $\beta$ 4GalT) [ 4-8] and  $\alpha$ 1,3 galactosyltransferase ( $\alpha$ 3GalT) genes [ 9, 10] have been identified. Similarly, four  $\beta$ 1,3 galactosyltransferase ( $\beta$ 3GalT) genes have been characterized recently [ 11-14].  $\beta$ 3GalT enzymes catalyze the formation of various structures of biological importance. For example, type 1 chain resulting from the transfer of galactose in a  $\beta$ 1,3 linkage to GlcNAc constitutes the base of the Gal $\beta$ 1,3(Fuc $\alpha$ 1,4)GlcNAc Lewis Le<sup>a</sup> and Fuc  $\alpha$ 1,2Gal $\beta$ 1,3(Fuc $\alpha$ 1,4)GlcNAc Le<sup>b</sup> antigens (reviewed in [ 15]). Also, galactose  $\beta$ 1,3 linked to GalNAc, a structure known as the Thomsen-Friedenreich antigen [ 16] enables

[List of Issues](#)

[Table of Contents](#)

[Prev Article](#)
[Next Article](#)

[Add to Favorite Articles](#)

[Email this to a Friend](#)



## QuickSearch in:

- ☒ Synergy  
☐ PubMed (MEDLINE)  
☐ CrossRef  
 for

## Authors:

- ☐ Dapeng Zhou  
☐ Eric G. Berger  
☐ Thierry Hennet  
☐

## Keywords

- ☐ galactosyltransferase  
☐ mucin  
☐ colon carcinoma  
☐ baculovirus  
☐

[Search](#)

(Received 19 February 1999, revised 4 May 1999, accepted 7 May 1999)

## Affiliations

Institute of Physiology, University of Zürich, Switzerland

## Correspondence

Correspondence to T. Hennet, Institute of Physiology, University of Zurich, Winterthurerstrasse

the formation of *core*2 structures of O-glycans [ 17]. Among the known  $\beta$ 3GalT enzymes, three catalyze the transfer of Gal to GlcNAc-based acceptors. Here, we report the characterization of an additional  $\beta$ 3GalT enzyme by searching for sequences similar to  $\beta$ 3GalT-I to -III proteins. The novel  $\beta$ 3GalT enzyme, named  $\beta$ 3GalT-V, displayed a marked preference for the O-linked *core*3 GlcNAc $\beta$ 1,3GalNAc acceptor structure. Accordingly, the  $\beta$ 3GalT-V enzyme likely represents the  $\beta$ 3GalT previously isolated from pig trachea [ 18].

190, 8057 Zurich, Switzerland.  
Fax: + 41 1 635 6814;  
Tel.: + 41 1 635 5080, E-mail: thennet@access.unizh.ch

## Materials and methods

Go to:



## Cloning of the $\beta$ 3GalT-V gene and baculovirus expression

The protein-coding region of the  $\beta$ 3GalT-V gene was amplified by PCR from human genomic DNA using the upstream primer 5'-GATTTGGATCCTTTCAGATGGCTTTC-3' and the downstream primer 5'-TGTGCCTCGAGGCTCCCTCAGACAG-3' that included *Bam*HI and *Xho*I restriction sites, respectively. The PCR conditions were three cycles at 94 °C for 45 s, 50 °C for 30 s and 72 °C for 60 s, followed by 25 cycles at 94 °C for 45 s, 58 °C for 30 s, 72 °C for 60 s. The resulting 956-bp fragment was subcloned into the *Bam*HI-*Xho*I sites of the pFastBac1 donor vector (BAC-to-BAC system, Life Technologies), which was subsequently processed as recommended by the manufacturer to generate pure recombinant baculoviruses. Sf9 insect cells were infected at a multiplicity of 10 and incubated at 27 °C for 72 h.

## Galactosyltransferase assay

The carbohydrate acceptors and the donor UDP-Gal were purchased from Sigma. Sf9 cells were lysed in 2% Triton X-100 for 15 min on ice and enucleated by centrifugation at 500 *g*. Galactosyltransferase activity was assessed using 10  $\mu$ L of Sf9 cell lysate in 50  $\mu$ L assays of 50 m M cacodylate buffer, pH 7.0, 10 m M MnCl<sub>2</sub>, 0.2 m M UDP-Gal,  $5 \times 10^4$  c.p.m. of UDP-[<sup>14</sup>C]Gal (82 pmol) (Amersham), and various acceptors ( Table 1). Samples were incubated at 37 °C for 60 min and reactions were stopped by adding 500  $\mu$ L of cold H<sub>2</sub>O. For kinetic analysis, the assays were performed using 0.5 m M UDP-Gal and  $10^5$  c.p.m. of UDP-[<sup>14</sup>C]Gal (164 pmol); reactions were stopped after 30 min. Samples were purified on Sep-Pak C<sub>18</sub> cartridges (Waters) by washing with 15 mL of H<sub>2</sub>O and eluting with 5 mL methanol. Samples containing ovalbumin as acceptor were precipitated with 1 mL of cold 15% trichloroacetic acid and 5% phosphotungstic acid solution and treated as described previously [ 19]. The amount of [<sup>14</sup>C]Gal transferred to the acceptors was measured in a liquid scintillation  $\beta$ -counter (Rackbeta, Pharmacia).


## Galactosidase assay

The disaccharide standards [<sup>14</sup>C]Gal $\beta$ 1,3GlcNAc- $\beta$ -O-p-nitrophenol ( $\beta$ -OpNP) and [<sup>14</sup>C]Gal $\beta$ 1,4GlcNAc( $\beta$ -OpNP) were synthesized using the  $\beta$ 3GalT-I [ 11] and  $\beta$ 4GalT [ 19] enzymes. Approximately 20 pmol of the purified standards and 20 pmol of [<sup>14</sup>C]-labeled  $\beta$ 3GalT-V reaction product were digested for 24 h at 37 °C with 2 U of recombinant *Xanthomonas manihotis*  $\alpha$ -galactosidase (New England Biolabs Inc.) in 50 m M sodium citrate, pH 6.5, 5 m M CaCl<sub>2</sub> and with 2 U of recombinant *X. manihotis*  $\beta$ -galactosidase (New England Biolabs Inc.) in 50 m M sodium citrate, pH 4.5, respectively. The reaction products were purified on Sep-Pak C<sub>18</sub> cartridges and the amount of [<sup>14</sup>C]Gal remaining bound to the acceptors was measured in the  $\beta$ -counter.

## Northern blotting


Total RNA from SK-N-MC, HeLa, HL-60, A-431, Caco2, Colo205 and PANC-1 cell lines was isolated by the procedure of Chomczynski and Sacchi [ 20]. Five micrograms of each tissue RNA were separated on 1% agarose-formaldehyde gels. Gels were rinsed in diethyl pyrocarbonate -treated water for 1 h and either stained in ethidium bromide, 0.1 M ammonium acetate or transferred to Hybond-N-Plus membranes (Amersham).  $\beta$ 3GalT mRNA in human tissues was detected by Northern blot analysis using commercially available multiple tissues poly(A)<sup>+</sup> RNA blots (Clontech). A *Bam*HI-*Xho*I

## Image Previews



[Full Size]

Table 1. Acceptor substrate specificity of the  $\beta$ 3GalT-I and  $\beta$ 3GalT-V enzymes.





956-bp fragment from the  $\beta$ 3GalT-V gene was labeled with [ $\alpha$ - $^{32}$ P]CTP (Hartmann Analytics, Braunschweig, Germany) by random priming. Membranes were hybridized for 16 h at 68 °C according to the manufacturer, then washed in 0.1  $\times$  NaCl/Cit, 0.1% SDS up to 55 °C and exposed for three days on Biomax-MS film (Kodak) using the appropriate intensifying screens.

## Results

Go to:




The protein sequences of the  $\beta$ 3GalT-I, -II and -III genes were compared to the EST division of GenBank using BLAST search algorithms [ 21]. The 338-bp human EST AJ003597 contained an open reading frame that was significantly similar to the three query sequences. The gene encoding the AJ003597 EST was identified within the PAC clone 70I24 (AF064860), which corresponds to a 165-kb stretch of chromosome 21 mapping to position 21q22.3. This gene included an open reading frame of 933 bp, which encoded a protein of 310 amino acids that was tentatively named  $\beta$ 3GalT-V (GenBank accession number AF145784). The  $\beta$ 3GalT-V protein contained a single membrane-spanning domain as deduced by Kyte-Doolittle hydropathy analysis [ 22]. The putative transmembrane domain began at amino acid position 8 and was 21 amino acids long, thus supporting a type-II transmembrane topology typical of most glycosyltransferases [ 3]. Phylogenetic analysis of the human galactosyltransferase enzymes indicated that  $\beta$ 3GalT-V was most closely related to the  $\beta$ 3GalT proteins, whereas these genes probably diverged consecutively to gene duplication events ( Fig. 1).

The  $\beta$ 3GalT-V protein shared 34%, 27%, 31% and 23% sequence identity with  $\beta$ 3GalT-I, -II, -III and -IV enzymes, respectively. The comparison with the  $\beta$ 3GalT-I, -II, -III proteins, which were used as the query for the retrieval of  $\beta$ 3GalT-V, indicated several conserved regions within the C-terminal two-thirds of the protein sequence, which corresponds to the catalytic domain according to the typical organization of glycosyltransferases ( Fig. 2). As expected, the N-terminal third encoding the cytoplasmic, transmembrane and stem domains, did not show much similarity to the other  $\beta$ 3GalT enzymes. The  $\beta$ 3GalT-V enzyme included all sequence motifs conserved among  $\beta$ 3GalT proteins [ 11-14]. In addition, six cysteine residues, which are conserved in  $\beta$ 3GalT-I to -III enzymes but absent in the structurally related  $\beta$ 3GalT-IV and  $\beta$ 3GnT [ 23] enzymes, are also found in the  $\beta$ 3GalT-V amino acid sequence, indicating that this enzyme is mostly related to the group of  $\beta$ 3GalTs that transfer Gal to GlcNAc-based acceptors.


To confirm the galactosyltransferase activity of the cloned gene, we constructed a recombinant baculovirus including the human  $\beta$ 3GalT-V gene by applying a transposon-mediated recombination system [ 24]. We have compared the galactosyltransferase activity in the lysate of Sf9 insect cells infected with recombinant baculoviruses expressing the  $\beta$ 3GalT-V and the  $\beta$ 3GalT-I enzyme [ 11]. Both  $\beta$ 3GalT enzymes displayed a preference towards GlcNAc-based acceptors and failed to transfer Gal to GalNAc ( Table 1). However,  $\beta$ 3GalT-V was able to transfer Gal to the O-linked core3 substrate whereas  $\beta$ 3GalT-I did not use this disaccharide acceptor efficiently. Another difference between both enzymes was detected when using the N-linked glycoprotein ovalbumin as an acceptor substrate, as  $\beta$ 3GalT-V catalyzed only a small incorporation of Gal onto this acceptor protein. Assays performed using UDP-GalNAc and UDP-GlcNAc as donors revealed that the two enzymes exhibit neither N-acetylglucosaminyltransferase nor N-acetylgalactosaminyltransferase activity (data not shown). As  $\beta$ 3GalT-V was able to transfer Gal to GlcNAc( $\beta$ -OpNP) and GlcNAc  $\beta$ 1,3GalNAc( $\alpha$ -OpNP), we have determined the  $K_m$  of the enzyme for these two acceptors. Saturation curves yielded apparent  $K_m$  values of 0.82 m M and 2.09 m M for GlcNAc  $\beta$ 1,3GalNAc( $\alpha$ -OpNP) and GlcNAc( $\beta$ -OpNP), respectively, thereby indicating the preference of  $\beta$ 3GalT-V for the O-linked core3 acceptor substrate ( Fig. 3).

The type of glycosidic linkage catalyzed by the  $\beta$ 3GalT-V enzyme was analyzed by *X. manihotis* galactosidase digestion [ 25]. The product of the reaction catalyzed by  $\beta$ 3GalT-V using the acceptor GlcNAc( $\beta$ -OpNP) and the donor UDP-Gal was purified on




[Full Size]

Fig. 3. Kinetic analysis of  $\beta$ 3GalT V. The  $K_m$  values for the acceptors GlcNAc $\beta$ 1,3GalNAc( $\alpha$ -OpNP) (A) and Glc...



[Full Size]

Fig. 4. Galactosidase digestion of the  $\beta$ 3GalT-V reaction product. The [ $^{14}$ C]Gal  $\beta$ 1,3GlcNAc( $\beta$ -OpNP) and [ $^{14}$ C]...




[Full Size]

Fig. 5. Expression pattern of the  $\beta$ 3GalT-V gene in adult human tissues. Each lane represents about 2  $\mu$ g o...

Sep-Pak C<sub>18</sub> cartridges and digested with galactosidases. Like the disaccharide Gal  $\beta$ 1,3GlcNAc ( $\beta$ -OpNP), the  $\beta$ 3GalT-V reaction product was cleaved by the *X. manihotis*  $\beta$ -galactosidase whereas it was found to be insensitive to  $\alpha$ -galactosidase treatment (Fig. 4). In comparison, the disaccharide Gal  $\beta$ 1,4GlcNAc ( $\beta$ -OpNP) remained intact after  $\beta$ -galactosidase digestion, thus showing the specificity of this galactosidase towards the  $\beta$ 1,3 linkage (Fig. 4).

The expression of the  $\beta$ 3GalT-V gene was investigated in adult human tissues by Northern blot analysis.  $\beta$ 3GalT-V transcripts were detected in small intestine, pancreas and testis (Fig. 5). Two transcripts of 7 kb and of  $\geq$  10 kb were visible in small intestine. Only the mRNA larger than 10 kb was detected in pancreas and testis, although other mRNAs may be present below the detection limit considering the weak hybridization signal observed in those tissues. By comparison, a large transcript of 7 kb was also detected for the  $\beta$ 3GalT-I gene [11, 14]. The expression pattern of the  $\beta$ 3GalT-V gene was also analyzed in tumor cell lines of different tissue origin.  $\beta$ 3GalT-V transcripts were visible in the pancreatic adenocarcinoma PANC-1 line and in the colorectal adenocarcinoma Colo205 cells (Fig. 6). By contrast,  $\beta$ 3GalT-V mRNA was not detectable in another colorectal adenocarcinoma line, Caco2, and in neuroblastoma SK-N-MC, squamous carcinoma A-431, promyelocytic leukemia HL-60 and cervical carcinoma HeLa cells.




[Full Size]

Fig. 6. Expression of the  $\beta$ 3GalT-V gene in tumor cells. Total RNA (5  $\mu$ g) from tumor cell lines was separa...

## Discussion

Go to:



The recent characterization of  $\beta$ 3GalT genes has underlined the structural differences between these genes and the other members of the galactosyltransferase family. It appears that  $\beta$ 3GalT enzymes share conserved sequences that are not found in  $\beta$ 4GalT and  $\alpha$ 3GalT proteins. These unique structural features have been exploited to identify additional  $\beta$ 3GalT genes, as illustrated in the present report describing the cloning of the human  $\beta$ 3GalT-V. Besides the amino acid sequence conservation, the  $\beta$ 3GalT-V gene also displayed a similar genomic organization, i.e. the entire protein-coding sequence is included in a single exon, as deduced by examination of the genomic PAC clone 70I24. In addition, as observed for  $\beta$ 3GalT-I, the  $\beta$ 3GalT-V transcripts were unexpectedly large for an open reading frame of 0.9 kb.

The same enzyme as the one reported here has been recently described by others under the name  $\beta$ 3Gal-T5 [26]. That work showed the capability of the  $\beta$ 3GalT-V to direct the expression of type 1 Lewis antigens when transfected in Namalwa lymphoma cells. The activity of the  $\beta$ 3GalT-V enzyme towards GlcNAc was confirmed by our experiments based on the recombinant expression of  $\beta$ 3GalT-V in insect cells. In addition, the expression of recombinant  $\beta$ 3GalT-V revealed the preference of the enzyme for the O-linked core3 GlcNAc  $\beta$ 1,3GalNAc acceptor.  $\beta$ 3GalT-V was the only  $\beta$ 3GalT enzyme identified to date capable of efficiently transferring Gal to this acceptor [11]. Also, the poor activity of  $\beta$ 3GalT-V towards the N-linked glycoprotein ovalbumin supported the preference of this enzyme for mucin-type glycoprotein acceptors. These findings suggest that  $\beta$ 3GalT-V probably represents the same enzyme as the  $\beta$ 3GalT previously isolated from pig trachea [18].  $\beta$ 3GalT-V transcripts were not detected in lung tissue as analyzed by Northern blotting. However, this may indicate that the  $\beta$ 3GalT-V gene expression is restricted to tracheal tissue, which was not represented in the source of lung RNA analyzed.

The  $\beta$ 3GalT-V gene was expressed in small intestine, pancreas and testis but not in differentiated colon tissue, whereas  $\beta$ 3GalT-V transcripts were detected in colonic adenocarcinoma Colo205 cells. A  $\beta$ 3GalT activity has been previously described in Colo205 cells [27-29] and in normal colonic mucosa [30]. The work of Isshiki *et al.* [26] demonstrated that this  $\beta$ 3GalT activity, encoded by the  $\beta$ 3GalT-V enzyme, was directed towards lacto-series structures, thereby contributing to the formation of type 1 carbohydrate chains. However, the absence of  $\beta$ 3GalT-V transcript in Caco2 cells and in other colorectal cancer cell lines [26] indicates that the neo-expression of this gene is not a general feature of colonic adenocarcinomas. Caco2 and Colo205 cell lines will be useful in investigating the relationship between the expression of  $\beta$ 3GalT-V and the

## To cite this article

Zhou, Dapeng, Berger, Eric G. & Hennet, Thierry (1999)  
Molecular cloning of a human  
UDP-galactose:GlcNAc  
 $\beta$ 1,3GalNAc  $\beta$ 1,3  
galactosyltransferase gene  
encoding an O-linked core3-  
elongation enzyme.  
*European Journal of  
Biochemistry* 263 (2), 571-576.  
doi: 10.1046/  
j.1432-1327.1999.00541.x

changes in levels of type 1 chain carbohydrate structures such as Le<sup>a</sup> and Le<sup>b</sup> antigens occurring in colonic cancer.

## Acknowledgements

Go to:






















We thank Bea Berger for technical assistance. This work was supported by a Roche Research Foundation grant and by the Swiss National Science Foundation Grant 3100-046836.96.

## References

Go to:



1. Costache, M., Apoil, P.A., Cailleau, A., Elmgren, A., Larson, G., Henry, S., Blancher, A., Iordachescu, D., Oriol, R., Mollicone, R. (1997) Evolution of fucosyltransferase genes in vertebrates. *J. Biol. Chem.* **272**, 29721-29728.  
  
2. Clausen, H. & Bennett, E.P. (1996) A family of UDP-GalNAc-polypeptide N-acetylgalactosaminyl-transferases control the initiation of mucin-type O-linked glycosylation. *Glycobiology* **6**, 635-646.  
 
3. Kleene, R. & Berger, E.G. (1993) The molecular biology of glycosyltransferases. *Biochim. Biophys. Acta* **1154**, 283-325.  
 
4. Shaper, N.L., Shaper, J.H., Meuth, J.L., Fox, J.L., Chang, H., Kirsch, I.R., Hollis, G.F. (1986) Bovine galactosyltransferase: identification of a clone by direct immunological screening of a cDNA expression library. *Proc. Natl Acad. Sci. USA* **83**, 1573-1577.  

5. Lo, N.W., Shaper, J.H., Pevsner, J., Shaper, N.L. (1998) The expanding  $\beta$ 4-galactosyltransferase gene family: messages from the databanks. *Glycobiology* **8**, 517-526.  
  
6. Almeida, R., Amado, M., David, L., Lavery, S.B., Holmes, E.H., Merckx, G., Vankessel, A.G., Rygaard, E., Hassan, H., Bennett, E., Clausen, H. (1997) A family of human  $\beta$ 4-galactosyltransferases - cloning and expression of two novel UDP-galactose:  $\beta$ -N-acetylglucosamine  $\beta$ 1,4-galactosyltransferases,  $\beta$ 4Gal-T2 and  $\beta$ 4Gal-T3. *J. Biol. Chem.* **272**, 31979-31991.  
  
7. Sato, T., Furukawa, K., Bakker, H., van den Eijnden, D.H., Van Die, I. (1998) Molecular cloning of a human cDNA encoding  $\beta$ 1,4-galactosyltransferase with 37% identity to mammalian UDP-Gal: GlcNAc  $\beta$ 1,4-galactosyltransferase. *Proc. Natl Acad. Sci. USA* **95**, 472-477.  
 
8. Schwientek, T., Almeida, R., Lavery, S.B., Holmes, E.H., Bennett, E., Clausen, H. (1998) Cloning of a novel member of the UDP-galactose- $\beta$ -N-acetylglucosamine  $\beta$ 1,4-galactosyltransferase family,  $\beta$ 4Gal-T4, involved in glycosphingolipid biosynthesis. *J. Biol. Chem.* **273**, 29331-29340.  
  
9. Yamamoto, F., Marken, J., Tsuji, T., White, T., Clausen, H., Hakomori, S.-I. (1990) Cloning and characterization of DNA complementary to human UDP-GalNAc: Fuc  $\alpha$ 1,2Gal  $\alpha$ 1,3GalNAc transferase (histo-blood group A transferase) mRNA. *J. Biol. Chem.* **265**, 1146-1151.

MEDLINE

ISI Abstract

10. Joziassse, D.H., Shaper, J.H., van den Eijnden, D.H., Van Tunen, A.J., Shaper, N.L. (1989) Bovine  $\alpha$ 1-3-galactosyltransferase: isolation and characterization of a cDNA clone. identification of homologous sequences in human genomic DNA. *J. Biol. Chem.* **264**, 14290-14297.

MEDLINE

ISI Abstract

11. Hennet, T., Dinter, A., Kuhnert, P., Mattu, T.S., Rudd, P.M., Berger, E.G. (1998) Genomic cloning and expression of three murine UDP-galactose:  $\beta$ -N-acetylglucosamine  $\beta$ 1,3-galactosyltransferase genes. *J. Biol. Chem.* **273**, 58-65.



MEDLINE

ISI Abstract

12. Kolbinger, F., Streiff, M.B., Katopodis, A.G. (1998) Cloning of a human UDP-galactose: 2-acetamido-2-deoxy- D-glucose 3 $\beta$ -galactosyltransferase catalyzing the formation of type 1 chains. *J. Biol. Chem.* **273**, 433-440.



MEDLINE

ISI Abstract

13. Miyazaki, H., Fukumoto, S., Okada, M., Hasegawa, T., Furukawa, K. (1997) Expression cloning of rat cDNA encoding UDP-galactose: G<sub>D2</sub>  $\beta$ 1, 3-galactosyltransferase that determines the expression of G<sub>D1b</sub>/G<sub>M1</sub>/G<sub>A1</sub>. *J. Biol. Chem.* **272**, 24794-24799.



MEDLINE

ISI Abstract

14. Amado, M., Almeida, R., Carneiro, F., Lavery, S.B., Holmes, E.H., Nomoto, M., Hollingsworth, M.A., Hassan, H., Schwientek, T., Nielsen, P.A., Bennett, E.P., Clausen, H. (1998) A family of human  $\beta$ -3-galactosyltransferases - Characterization of four members of a UDP-galactose- $\beta$ -N-acetylglucosamine/ $\beta$ -N-acetylgalactosamine  $\beta$ -1,3-galactosyltransferase family. *J. Biol. Chem.* **273**, 12770-12778.



MEDLINE

ISI Abstract

15. Lowe, J.B. (1995) Biochemistry and biosynthesis of ABH and Lewis antigens. In *Molecular Basis of Human Blood Group Antigens* (Cartron, J.-P. & Rouger, P., eds), pp. 75-115. Plenum Press, New York.

16. Thomas, D.B. & Winzler, R.J. (1969) Structural studies on human erythrocyte glycoproteins. Alkali-labile oligosaccharides. *J. Biol. Chem.* **244**, 5943-5946.

MEDLINE

ISI Abstract

17. Brockhausen, I. (1995) Biosynthesis of O-glycans of the N-acetyl galactosamine- $\alpha$ -Ser/Thr linkage type. In *Glycoproteins* (Montreuil, J., Vliegenthart, J.F.G. & Schachter, H., eds), pp. 201-259. Elsevier, Amsterdam.

18. Sheares, B.T. & Carlson, D.M. (1983) Characterization of UDP-galactose: 2-acetamido-2-deoxy- D-glucose 3 $\beta$ -galactosyltransferase from pig trachea. *J. Biol. Chem.* **258**, 9893-9898.

MEDLINE

ISI Abstract

19. Malissard, M., Borsig, L., Di Marco, S., Grutter, M.G., Kragl, U., Wandrey, C., Berger, E.G. (1996) Recombinant soluble  $\beta$ -1,4-galactosyltransferases expressed in *Saccharomyces cerevisiae*. Purification, characterization and comparison with human enzyme. *Eur. J. Biochem.* **239**, 340-348.

MEDLINE

ISI Abstract

20. Chomczynski, P. & Sacchi, N. (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* **162**, 156-159.





























MEDLINE

ISI Abstract

21. Altschul, S.F., Boguski, M.S., Gish, W., Wootton, J.C. (1994) Issues in searching molecular sequence databases. *Nat. Genet.* **6**, 119-129.

MEDLINE

ISI Abstract

22. Kyte, J. & Doolittle, R.F. (1982) A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**, 105 132.  
  
23. Zhou, D., Dinter, A., Gutierrez Gallego, R., Kamerling, J.P., Vliegenthart, J.F.G., Berger, E.G., Hennet, T. (1999) A  $\beta$ -1,3-*N*-acetylglucosaminyltransferase with poly-*N*-acetyllactosamine synthase activity is structurally related to  $\beta$ -1,3-galactosyltransferases. *Proc. Natl Acad. Sci. USA* **96**, 406 411.  
 
24. Luckow, V.A., Lee, S.C., Barry, G.F., Olins, P.O. (1993) Efficient generation of infectious recombinant baculoviruses by site-specific transposon-mediated insertion of foreign genes into a baculovirus genome propagated in *Escherichia coli*. *J. Virol.* **67**, 4566 4579.  
  
25. Wong-Madden, S.T. & Landry, D. (1995) Purification and characterization of novel glycosidases from the bacterial genus *Xanthomonas*. *Glycobiology* **5**, 19 28.  

26. Isshiki, S., Togayachi, A., Kudo, T., Nishihara, S., Watanabe, M., Kubota, T., Kitajima, M., Shiraishi, N., Sasaki, K., Andoh, T., Narimatsu, H. (1999) Cloning, expression, and characterization of a novel UDP-galactose:  $\beta$ -*N*-acetylglucosamine  $\beta$ 1,3-galactosyltransferase ( $\beta$ 3Gal-T5) responsible for synthesis of type 1 chain in colorectal and pancreatic epithelia and tumor cells derived therefrom. *J. Biol. Chem.* **274**, 12499 12507.  
  
27. Holmes, E.H. & Levery, S.B. (1989) Preparative in vitro generation of lacto-series type 1 chain glycolipids catalyzed by  $\beta$ 1,3-galactosyltransferase from human colonic adenocarcinoma colo 205 cells. *Arch. Biochem. Biophys.* **274**, 14 25.  
 
28. Holmes, E.H. (1989) Characterization and membrane organization of  $\beta$ 1,3- and  $\beta$ 1, 4-galactosyltransferases from human colonic adenocarcinoma cell lines Colo 205 and SW403: basis for preferential synthesis of type 1 chain lacto-series carbohydrate structures. *Arch. Biochem. Biophys.* **270**, 630 646.  
  
29. Valli, M., Gallanti, A., Bozzaro, S., Trinchera, M. (1998)  $\beta$ -1,3-Galactosyltransferase and  $\beta$ -1,2-fucosyltransferase involved in the biosynthesis of type-1-chain carbohydrate antigens in human colon adenocarcinoma cell lines. *Eur. J. Biochem.* **256**, 494 501.  
  
30. Seko, A., Ohkura, T., Kitamura, H., Yonezawa, S., Sato, E., Yamashita, K. (1996) Quantitative differences in GlcNAc:  $\beta$ 1 $\rightarrow$ 3 and GlcNAc:  $\beta$ 1 $\rightarrow$ 4 galactosyltransferase activities between human colonic adenocarcinomas and normal colonic mucosa. *Cancer Res.* **56**, 3468 3473.  
  
31. Page, R.D.M. (1996) TREEVIEW: an application to display phylogenetic trees on personal computers. *Computer Applic. Biosci.* **12**, 357 358.  

32. Thompson, J.D., Higgins, D.G., Gibson, T.J. (1994) Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673 4680.  
 

## Abbreviations

Go to:  $\beta$ 3GalT  $\beta$ 1,3 galactosyltransferase $\beta$ 4GalT  $\beta$ 1,4 galactosyltransferase $\alpha$ 3GalT  $\alpha$ 1,3 galactosyltransferase

Bn benzyl

pNP *p*-nitrophenol

EST expressed sequence tag.

Forward Links to Citing  
ArticlesGo to: 

- Lydia Mare and Marco Trinchera . Suppression of  $\beta$ 1,3galactosyltransferase  $\beta$ 3Gal-T5 in cancer cells reduces sialyl-Lewis a and enhances poly N-acetyllactosamines and sialyl-Lewis x on O-glycans. *European Journal of Biochemistry* 271: 1, 186-194.

[Abstract](#) [References](#) [Full Text Article](#) [PDF](#)
- Martine Malissard, André Dinter, Eric G. Berger and Thierry Hennet . Functional assignment of motifs conserved in  $\beta$ 1,3-glycosyltransferases. A mutagenesis study of murine UDP-galactose: $\beta$ -N-acetylglucosamine  $\beta$ 1,3-galactosyltransferase-I. *European Journal of Biochemistry* 269: 1, 233-239.

[Abstract](#) [References](#) [Full Text Article](#) [PDF](#)

## Footnotes

Note: the nucleotide sequence reported in this paper has been submitted to the GenBank™/EBI Data Bank with accession number AF145784.

## European Journal of Biochemistry

Volume 263 Issue 2 Page 571 - July (II) 1999

Blackwell Synergy® is a Blackwell Publishing, Inc. registered trademark  
 More information about Blackwell Synergy - online journals from [www.blackwellpublishing.com](http://www.blackwellpublishing.com).  
 We welcome your Feedback. See our [Privacy Statement](#) and [Terms and Conditions](#).  
 Technology Partner - [Atypion Systems, Inc.](#)

